



Wiadomości Lekarskie

Official journal of the Polish Medical Association



Memory of
dr Władysław
Biegański

VOLUME LXXIV, ISSUE 9 PART 1, SEPTEMBER 2021

Since 1928



ALUNA Publishing House

Wiadomości Lekarskie is abstracted and indexed in: PUBMED/MEDLINE, SCOPUS, EMBASE, INDEX COPERNICUS, POLISH MINISTRY OF EDUCATION AND SCIENCE, POLISH MEDICAL BIBLIOGRAPHY

Copyright: © ALUNA Publishing House.

Articles published on-line and available in open access are published under Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Wiadomości Lekarskie monthly journal

You can order the subscription for the journal from Wydawnictwo Aluna by:

prenumerata@wydawnictwo-aluna.pl
Wydawnictwo Aluna
Z.M. Przesmyckiego 29
05-510 Konstancin-Jeziorna
Poland

Place a written order first.

If you need, ask for an invoice.

Payment should be done to the following account of the Publisher:

account number for Polish customers (PLN):

82 1940 1076 3010 7407 0000 0000

Credit Agricole Bank Polska S. A., SWIFT: AGRIPPLR

account number for foreign customers (EURO):

57 2490 0005 0000 4600 7604 3035

Alior Bank S. A.: SWIFT: ALBPPLPW

Subscription of twelve consecutive issues (1-12):

Customers in Poland: 360 PLN/year

Customers from other countries: 320 EURO/year

Valentyna Sklyarova, Kateryna Shatylovich, Pavlo Sklyarov, Anzhelika Filipyuk SHOULD ASCARIASIS BE CONSIDERED AS A REPRODUCTOLOGY PROBLEM?	2138
Oleksandr Belov, Yevhen Loiko, Iryna Mazaikina, Iryna Pototska FEATURES OF PSYCHOPATHOLOGICAL REACTION IN PARENTS WHOSE CHILDREN SUFFER FROM CHRONIC MUSCULOSKELETAL DISORDERS	2147
Marta M. Zelinka-Khobzey, Kostiantyn V. Tarasenko, Tetiana V. Mamontova, Oksana A. Shlykova CHARACTERISTICS OF CD68+ AND CD163+ EXPRESSION IN PLACENTA OF WOMEN WITH PREECLAMPSIA AND OBESITY	2152
Yaroslav P. Feleshtynskiy, Sergiy O. Oparin, Bogdan V. Sorokin, Margaryta G. Boiarskaia, Dmytro V. Lutsenko ENDOSCOPIC HEMOSTASIS IN ULCERATIVE GASTRODUODENAL BLEEDING USING HIGH-FREQUENCY BIOLOGICAL WELDING ELECTROLIGATION	2159
Oksana V. Tymoshchuk, Zhanna Sotnikova-Meleshkina, Olha P. Melnyk-Sheremeta, Oksana V. Zahorodnieva, Krystyna A. Kryvonos, Kseniia M. Ostrovska KEY ASPECTS OF ORGANIZATION OF THE PROCESS AND INDIVIDUAL CHARACTERISTICS OF NUTRITION OF PUPILS AND STUDENTS OF DIFFERENT TYPES OF MODERN EDUCATIONAL INSTITUTIONS	2163
Nataliya O. Gutorova, Vitalii M. Pashkov, Oleksii S. Soloviov ILLEGAL INTERNET PHARMACIES AS A THREAT TO PUBLIC HEALTH IN EUROPE	2169
Natalya A. Maruta, Marianna V. Markova, Hanna M. Kozhyna, Tetiana A. Aliieva, Lyudmyla M. Yuryeva, Tsira B. Abdryakhimova, Nataliia G. Pshuk, Andrii M. Skrypnykov PSYCHOLOGICAL FACTORS AND CONSEQUENCES OF PSYCHOSOCIAL STRESS DURING THE PANDEMIC	2175
Igor S. Brodetskiy, Vladislav A. Malanchuk, Olena O. Dyadyk, Yaroslava A. Kulbashna, Olena O. Astapenko, Nataliia V. Kapustnyk, Mykhailo S. Myroshnychenko MORPHOLOGICAL JUSTIFICATION OF THE SALIVARY GLAND TISSUE RESECTION BOUNDARIES DURING SURGICAL TREATMENT OF PATIENTS WITH PLEOMORPHIC ADENOMAS	2182
Yurii Yu. Yarov, Iryna I. Tkachenko DYNAMICS OF ANTI-ANTIOXIDANT SYSTEM INDICATORS IN THE POSTOPERATIVE PERIOD IN PATIENTS WITH PERIODONTITIS ACCOMPANIED BY DIFFERENT REACTIVITY OF THE ORGANISM	2187
Vladislav V. Liubchak, Liliia M. Khomenko, Michael P. Kovalishyn, Viktoriia V. Ilyina, Tetiana V. Babar, Volodymyr V. Sikora, Olha V. Simonova NEW MARKETING TOOLS IN THE BLOOD SERVICE TO OPTIMIZE QUALITY MANAGEMENT	2192
Nazar R. Hrytsevych, Volodymyr V. Vereschaka FEATURES OF THE ULTRASTRUCTURE OF THE SKIN OF WHITE RATS 60 AND 90 DAYS AFTER MODELING OF PORTAL HYPERTENSION	2197
Roman Ye. Bulyk, Vladyslav R. Yosypenko, Kateryna V. Vlasova ONTOGENETIC ASPECTS OF MELATONIN RECEPTORS 1A DENSITY IN THE LATERAL PREOPTIC NUCLEUS OF THE HYPOTHALAMUS UNDER VARIOUS LIGHT CONDITIONS	2202
REVIEW ARTICLES	
Vitalii Pashkov, Maryna V. Trotska, Liudmyla M. Nikolenko AMBIENT AIR PROTECTION IN THE SYSTEM OF PUBLIC HEALTH AND BIOLOGICAL SECURITY LEGAL PROVISION	2207
Liudmyla A. Vygivska, Lesia A. Rudenko, Evgeniya B. Radzishavska, Iryna M. Merenkova, Violeta B. Kalnytska PECULIARITIES OF THE PREVALENCE OF INTRAUTERINE INFECTIONS AT THE PRESENT STAGE	2213
Anna Krakowiak, Marek Rajzer, Mateusz Gaczoł, Urszula Gancarczyk, Paweł Prochownik, Natalia Podolec, Zuzanna Sachajko, Filip Baranowski, Anna Pyczek, Monika Komar OBESITY AND ATRIAL FIBRILLATION – BARIATRIC SURGERY AS A METHOD OF AF RISK DECREASE	2218
Halyna V. Bilavych, Jan V. Bilawicz, Larysa V. Slyvka, Mariia V. Klepar, Mykola P. Pantiuk, Nadiya O. Fedchyshyn, Borys P. Savchuk SANITARY AND HYGIENIC EDUCATION OF CHILDREN AND ADULTS UNDER THE CONDITIONS OF THE COVID-19 PANDEMIC IN MEASUREMENT OF THE ACHIEVEMENTS OF PUBLIC MEDICINE OF GALICIA (END OF THE XIX – 30S OF THE XX CENTURY)	2222
Waldemar Gniadek MIECZYSŁAW KONOPACKI – THE OUTSTANDING POLISH HISTOLOGIST, ANATOMIST, AND FREEMASON ACTIVIST	2228
CASE STUDIES	
Vladimir Belokonev, Andrei I. Gritsaenko, Alexey Nikolaev, Tatiana Larina A CLINICAL CASE OF COEXISTENT PARASITIC LIVER AND GASTRIC LESIONS CAUSED BY CLONORCHIASIS	2232
Aleksandra Oraczewska, Marek Ochman, Mariola Ograbek-Król, Wiktoria Kowalska, Klaudia Glinka, Fryderyk Zawadzki, Adam Barczyk, Szymon Skoczyński PULMONARY ALVEOLAR MICROLITHIASIS. DISCREPANCIES BETWEEN RADIOLOGICAL FINDINGS AND CLINICAL PATTERN- CASE STUDY	2235

ENDOSCOPIC HEMOSTASIS IN ULCERATIVE GASTRODUODENAL BLEEDING USING HIGH-FREQUENCY BIOLOGICAL WELDING ELECTROLIGATION

DOI: 10.36740/WLek202109123

Yaroslav P. Feleshtynskyi¹, Sergiy O. Oparin², Bogdan V. Sorokin¹, Margaryta G. Boiarskaia³, Dmytro V. Lutsenko²¹SHUPYK NATIONAL HEALTHCARE UNIVERSITY OF UKRAINE OF THE MINISTRY OF HEALTH OF UKRAINE, KYIV, UKRAINE²MUNICIPAL NON-COMMERCIAL ENTERPRISE OF THE KYIV REGIONAL COUNCIL "KYIV REGIONAL CLINICAL HOSPITAL", KYIV, UKRAINE³KYIV CITY CLINICAL HOSPITAL No.12 (CITY CENTER FOR GASTROINTESTINAL BLEEDING), KYIV, UKRAINE

ABSTRACT

The aim: To increase the efficiency of endoscopic hemostasis in ulcerative gastroduodenal bleeding using high-frequency biological welding electroliation.**Materials and methods:** The evaluation of endoscopic hemostasis in 160 patients aged 40 to 85 years with ulcerative gastroduodenal bleeding for the period from 2017 to 2020 was carried out. The patients were divided into two groups: the first (treatment) group involved 80 patients who underwent high-frequency biological welding electroliation, the second (experimental) group consisted of 80 patients who underwent monopolar thermal argon plasma coagulation.**Results:** In the first (treatment) group of patients with ulcerative gastroduodenal bleeding, who underwent endoscopic hemostasis using high-frequency biological welding electroliation, primary hemostasis was achieved in 77 cases (96.25%). In the first group, an early recurrence of bleeding was registered in 3 patients (3.75%). In the second (experimental) group of patients with ulcerative gastroduodenal bleeding, who underwent endoscopic hemostasis using monopolar thermal argon plasma coagulation, primary hemostasis was achieved in 66 cases (82.5%). In the second group, a recurrence of bleeding was observed in 14 patients (17.5%).**Conclusions:** The use of high-frequency biological welding electroliation for endoscopic hemostasis in ulcerative gastroduodenal bleeding provides a more reliable permanent hemostasis compared to the use of monopolar thermal argon plasma coagulation (77 (96.5%) and 66 (82.5%) cases, respectively). The frequency of bleeding recurrence is reduced to 3.5% and 17.5%, respectively, and the number of surgical interventions for acute bleeding in case of recurrence is decreased to 3 (3.5%) and 7 (8.75%), respectively.**KEY WORDS:** ulcerative gastroduodenal bleeding, endoscopic probe, endoscopic hemostasis, high-frequency biological welding electroliation, argon plasma coagulation

Wiad Lek. 2021;74(9 p.1):2159-2162

INTRODUCTION

At present, ulcerative gastroduodenal bleeding remains a challenging and not fully resolved problem in view of the complexity of tactical approaches, poor treatment outcomes, high overall mortality rate, which is 8-14% [1,2]. Despite the success of endoscopic hemostasis, intensive therapy, the use of modern methods of surgical interventions, postoperative mortality ranges from 10 to 25% [1,2], and in case of severe ulcerative gastroduodenal bleeding it might be within the range of 15-50% [1,3], as well as 30-70% in case of recurrence [1,3].

The analysis of causes of poor treatment outcomes shows that the achievement of the desired results should involve the perfection of methods of endoscopic hemostasis and prevention of early recurrence of bleeding in order to reduce the mortality rate of patients with ulcerative gastroduodenal bleeding. A number of methods of endoscopic hemostasis are in wide use for ulcerative gastroduodenal bleeding control, including chemical (caproferrum, lifusolum), injection (epinephrine : terlipressin 1:20), thermal – monopolar coagulation (diathermocoagulation, hydrodiathermocoagulation, argon plasma coagulation, laser

photocoagulation), mechanical (clipping and ligation), combined (injection + argon plasma), etc. Despite the large number of existing methods of endoscopic hemostasis, each has both advantages and disadvantages, which are accompanied by a high recurrence rate of ulcerative gastroduodenal bleeding.

Currently, the most common method of endoscopic hemostasis is monopolar thermal argon plasma coagulation [4]. Despite all its advantages, it still gives cause for concern. A number of experts [4] still believe that argon plasma coagulation does not provide a sufficient coagulation effect on tissues and cannot reliably stop the bleeding from arterial vessels with a diameter of 2 mm or more deep in the tissue array, and there is a risk of damaging during the procedure of large-diameter vessels that do not bleed, especially in case of ulcerative gastroduodenal bleeding [5].

The literature indicates that the percentage of early recurrence in endoscopic hemostasis using argon plasma coagulation ranges from 15% to 22% [5-11]. In this regard, an alternative method of endoscopic hemostasis, namely high-frequency biological welding electroliation has been proposed.

THE AIM

To increase the efficiency of endoscopic hemostasis in ulcerative gastroduodenal bleeding using high-frequency biological welding electroligation.

MATERIALS AND METHODS

The evaluation of endoscopic hemostasis in 160 patients with ulcerative gastroduodenal bleeding for the period from 2017 to 2020 was carried out. The age of patients ranged from 40 to 85 years. The average age was 62.5 ± 9.1 years. There were 110 men and 50 women.

The patients were under medical treatment at the clinic of the Department of Proctology of Shupyk National Healthcare University of Ukraine.

Depending on the method of endoscopic hemostasis, the patients were divided into two groups: the first (treatment) group involved 80 patients who underwent high-frequency biological welding electroligation, the second (experimental) group consisted of 80 patients who underwent monopolar thermal argon plasma coagulation. In patients of the first group, endoscopic hemostasis was performed according to the developed technique (Patent of Ukraine for invention UA No.121365) [12].

According to the Forrest classification of bleeding [7], the patients were distributed as follows: the first (treatment) group: FIA – 3 patients (3.75%); FIB – 14 (17.5%); FIX – 4 (5%); FIIA – 28 (35%); FIIB – 20 (25%); FIIC – 11 (13.75%); the second (experimental) group: FIA – 7 patients (8.75%); FIB – 13 (16.25%); FIX – 8 (10%); FIIA – 28 (35%); FIIB – 19 (23.75%); FIIC – 5 (6.25%).

Among 80 patients of the treatment group with ulcerative gastroduodenal bleeding, there were 52 men (65%) and 28 women (35%). According to localization, there were 33 (40.25%) cases of gastric ulcers; 44 (56.75%) cases of duodenal ulcers, including 3 cases of large penetrating ulcers; and 3 (3.75%) cases of postbulbar ulcers. Among 80 patients of the experimental group with ulcerative gastroduodenal bleeding, there were 58 men (72.5%) and 22 women (27.5%). According to localization, there were 31 (38.75%) cases of gastric ulcers, including 3 cases of large penetrating ulcers; 46 (57.5%) cases of duodenal ulcers; and 3 (3.75%) cases of postbulbar ulcers. Thus, the studied groups are comparable in sex composition ($p=0.306$) and localization of bleeding ($p=0.948$).

The sources of gastrointestinal bleeding in 160 patients were the following: gastric ulcers (38.3%), duodenal ulcers (58%), postbulbar ulcers (3.70%). The diameter of bleeding ulcerous defects was from 5 mm to 40 mm ($12.2 \text{ mm} \pm 1.2$ on average).

The technique of endoscopic hemostasis of ulcerative gastroduodenal bleeding using high-frequency biological welding electroligation was performed on the EKVZ-300 Patonmed device in automatic welding mode – subtype 5. During videogastroduodenoscopy, a welding high-frequency bipolar probe with an electrode located at the end of the probe is passed through the forceps aperture of the video endoscope directly to the vessel, which bleeds. In

this case, the exposure is 15-40 seconds, depending on the vessel diameter. If the vessel diameter is 1-2 mm of a stomach or duodenum ulcer, a 2.8 mm probe with a concave electrode is used through a video gastroscope, with a power of 60-70 W and a temperature of 60 °C at the end of the electrode. If the vessel diameter is 2-3 mm of a postbulbar ulcer, a 3.2 mm probe with a concave electrode is used through a video duodenoscopy, with a power of 70-80 W and a temperature of 70-80 °C at the end of the electrode.

In patients of the second (experimental) group, endoscopic hemostasis was performed using monopolar thermal argon plasma coagulation on the Bowa device. Coagulation occurs without direct contact between the active electrode and the tissue, while the argon flow displaces oxygen from the coagulation zone, which significantly reduces the carbonization (burning) of the tissue. Coagulation was also performed under an angle, as the plasma jet was deflected towards the most conductive (wet) tissues, namely fresh blood, clots.

The evaluation of endoscopic hemostasis was carried out taking into account the effectiveness of hemostasis and the frequency of bleeding recurrence.

RESULTS

The result of endoscopic hemostasis in ulcerative gastroduodenal bleeding in the treatment group of patients who underwent welding electroligation, as well as in the experimental group of patients who underwent argon plasma coagulation are presented in table I.

In the first (treatment) group of patients with ulcerative gastroduodenal bleeding, who underwent endoscopic hemostasis using high-frequency biological welding electroligation, primary hemostasis was achieved in 77 cases (96.25%). In this study group, an early recurrence of bleeding after endoscopic hemostasis was registered in 3 patients (3.75%) in the first 24 hours. In the second (experimental) group of patients with ulcerative gastroduodenal bleeding, who underwent endoscopic hemostasis using monopolar thermal argon plasma coagulation, primary hemostasis was achieved in 66 cases (82.5%). In this study group, a recurrence of bleeding was observed in 14 patients (17.5%). This statistical evidence coincides with the literature data from various sources (15-22%).

In the main group, in 3 (3.75%) patients, endoscopic hemostasis was achieved by a combined method (injection + welding ligation). Combined endoscopic hemostasis was performed in FIA in 3 (3.75%) patients. These patients had large duodenal penetrating ulcers with bleeding vessels larger than 5 mm in diameter. As a supplementary method for combined endoscopic hemostasis, the injection method was used, in particular adrenaline solution (1:5000). Due to recurrence of bleeding and ineffectiveness of endoscopic hemostasis, 3 patients of the treatment group underwent surgery. They were discharged and referred to outpatient treatment in satisfactory condition.

In the second (experimental) group of patients with ulcerative gastroduodenal bleeding, who underwent

Table I. Comparative analysis of treatment outcomes.

Parameter	Treatment group N=80	Experimental group N=80	OR (95% CI)	p
Primary hemostasis	77 (96.25%)	66 (82.5%)	5.4 (1.4-30.5)	0.005*
Bleeding recurrence	3 (3.7%)	14 (17.5%)	0.18 (0.03-0.7)	0.005*

OR (95% CI) – relative risk assessment (odds ratio) when comparing the treatment and the experimental groups; p – statistical significance chi-squared test (Fisher's exact test* – $p < 0.05$).

endoscopic hemostasis using monopolar thermal argon plasma coagulation, endoscopic hemostasis was performed by a combined method (injection + APC) in 14 (17.5%) patients. On the first day, recurrence was registered in 5 (6.25%) patients, on the second and third days – in 9 (11.25%) patients. In these patients with recurrent bleeding, the vessel diameter was more than 2 mm.

During monopolar thermal argon plasma coagulation, in 1 patient with peripapillary duodenum ulcer complicated by intestinal bleeding (Forrest IA, post-hemorrhagic shock of stage I-II), there was a complication in the form of perforation of the anterior-lateral wall of the duodenum. The patient underwent surgery and was discharged and referred to outpatient treatment in satisfactory condition.

In 7 patients with recurrent ulcerative bleeding (on the first and second days) after argon plasma coagulation, welding electroligation was applied resulting in permanent hemostasis. In the other 7 patients with recurrent bleeding and ineffective endoscopic hemostasis, surgical interventions were performed on the third day, which were effective. The patients were discharged and referred to outpatient treatment in satisfactory condition.

The comparison of the results in the study groups indicates a significant increase in the probability of achieving hemostasis in the treatment group by 5.4 times (OR=5.4, $p=0.005$); a decrease in the risk of recurrent bleeding in the treatment group by 82% (OR=0.18, $p=0.005$) and a decrease in the risk of surgical correction of recurrence in the treatment group by 87% (OR=0.13, $p=0.03$).

All patients of both groups with ulcerative gastroduodenal bleeding who underwent endoscopic hemostasis, were prescribed antiulcer drug therapy using proton pump inhibitors and follow-up videogastroscopy in the postoperative period.

DISCUSSION

The findings of the use of endoscopic hemostasis in ulcerative gastroduodenal bleeding showed that the best results were obtained in patients of the treatment group using high-frequency biological welding electroligation. In particular, recurrences of ulcerative bleeding were registered only in 3 (3.5%) patients of the treatment group, while there were 14 (17.5%) cases in the experimental group. Permanent hemostasis was achieved in 77 (96.25%) patients of the treatment group and 66 (82.5%) patients of the experimental group.

Higher efficiency of endoscopic hemostasis in patients of the treatment group was achieved due to vessel sealing by

means of the impedance action of a modulated high-frequency current signal and thanks to an adaptive system for automatic control of the welding process through the contact of a special endoscopic probe with a concave electrode (Cu + Mo alloy) [6]. Morphological verification of the efficiency of high-frequency biological welding electroligation of bleeding vessels with a diameter of 1-5 mm was experimentally confirmed. It was found that vessel sealing is followed by total obliteration due to coagulation of the endothelium of the vessel intima, disorganization of the inner elastic membrane, dissection of the outer membrane, covering the adventitia of the vessel [9]. The intestinal mucosa around the sealing zone is not exposed to high temperatures, resulting in a lower percentage of complications [6].

Argon plasma coagulation does not provide a sufficient coagulation effect on tissues and cannot reliably stop the bleeding from arterial vessels deep in the tissue array. Moreover, there is a risk of damaging the walls during argon plasma coagulation [8]. In particular, among patients of the experimental group with ulcerative gastroduodenal bleeding after argon plasma coagulation, a complication such as perforation was registered in one (1.25%) patient who required surgery. In the treatment group of patients where endoscopic hemostasis was performed using high-frequency biological welding electroligation, no perforations were registered.

The method of influencing the bleeding source of high-frequency current during biological welding electroligation is the electrical effect of a modulated signal. And during argon plasma coagulation, there is a thermal effect with the release of a large amount of heat. In addition, the difficulty of energy dosing in the contact mode of argon plasma coagulation provokes the threat of burning the wall of the hollow organ [15], while during welding electroligation, energy dosing is guided by an adaptive system for automatic control of the welding process [6].

Thus, the proposed technique of high-frequency biological welding electroligation has significant advantages over thermal argon plasma coagulation. It provides permanent hemostasis of the vessels from 1 to 5 mm, which in turn can reduce the recurrence rate of bleeding and surgical interventions for acute bleeding.

CONCLUSIONS

The use of high-frequency biological welding electroligation for endoscopic hemostasis in ulcerative gastroduodenal bleeding provides a more reliable permanent hemo-

stasis compared to the use of monopolar thermal argon plasma coagulation (96.5% and 82.5%, respectively). The frequency of bleeding recurrence is reduced to 3.5% and 17.5%, respectively, and the number of surgical interventions for acute bleeding in case of recurrence is decreased to 3 (3.5%) and 7 (8.75%), respectively.

REFERENCES

- Fomin P.D., Shepetko E.M. Hirurgichne likuvannya virazkovih gastroduodenalnih krovotech ta korekciya postgemoragichnoi anemii gemopoetichnimi stovburovimi klitinami. Kyiv: «Feniks». 2011: 7p. (in Ukrainian)
- Vinokurov M.M., Kapitonova M.A. Taktika likuvannya hvorih z virazkovimi gastroduodenalnimi krovotechami. Hirurgiya. 2008;2: 33-36. (in Ukrainian)
- Lebedev N.V., Klimov A.E., Barhudarova T.V., Malkarov M.A. Taktika likuvannya hvorih z virazkovimi gastroduodenalnimi krovotechami. Visnik hirurgii. 2007;166: 76-79. (in Ukrainian)
- Fedorov E.D., Yudin O.I., Petrov D.U. et al. Effektivnost primeneniya endoskopicheskoy argonoplazmennoj koagulyacii v sravnenii s bipolyarnoy diatermokoagulyaciej pri yazvennyh gastroduodenalnyh krovotecheniyah. RZHGGK. 2008;18: 33. (in Russian)
- Bereznickij Y.S., Sulima V.P., Yaroshenko K.O. Vibir metodu endoskopichnogo gemostazu pri gostrih shlunkovo-kishkovih krovotechah. 2016: 08-111. (in Ukrainian)
- Paton B.Y., Ivanova O.N. Tkaninozberigayucha visokochastotna elektrovaryvalna hirurgiya. Atlas. Kyiv: «Naukova dumka» NAN Ukraïni. 2009.
- Unifikovaniy klinichnij protokol pervinnoi, vtorinnoi (specializovanoi) ta tretinnoi (visokospecializovanoi) medichnoi dopomogi. Shlunkovo-kishkovi krovotechi: 2016. (in Ukrainian)
- Shepetko E.M. Hirurgichne likuvannya uskladnenih yukstapapilyarnih virazok. Kyiv. Zhurnal «Raduga». 2012: 12-15. (in Ukrainian)
- Oparin S.O., Felishtinskij Y.P., Dyaduh O.A. Eksperimentalne obruntuvannya endoskopichnogo likuvannya shlunkovo-kishkovih krovotech za dopomogoyu visokochastotnogo zvaryvalnogo elektrokoagulyatora. Zhurnal «Morfologiya» 2018: 105-110. (in Ukrainian)
- Akin M., Alkan E., Tuna Y., Yalcinkaya T. Comparison of heater probe coagulation and argon plasma coagulation in the management of Mallory-Weiss tears and high-risk ulcer bleeding. 2017;18(1):35-38. doi: 10.1016/j.ajg.2017.01.006.
- Akin M., Alkan E., Tuna Y., Yalcinkaya T. Comparison of heater probe coagulation and argonplasma coagulation in the management of Mallory-Weiss tears and high-risk ulcer bleeding. Gastroenterol. 2017;8: 52-54.
- Patent of Ukraine for invention №121365 UA; IPC A61V 18/08 (2006.01) A61N 01/04 (2006.01) A61B 17/22 (2006.01) A61B17/94(2006.01). Method of endoscopic treatment of ulcer gastroduodenal bleeding. – No a2019 02633; app. 08.03.19; published 12.05.20.
- Patent of Ukraine for utility model №115147 UA; IPC A61B 17/94 (2006.01) A61N1/18 (2006.01) High-frequency bipolar probe for endoscopic minimally invasive surgery. U2016 08620; app. 05.08.2016; publ. 10.04.2017.
- Patent of Ukraine for utility model №124885 UA; IPC A61B 18/08 (2006.01) A61B 17/94 (2006.01) A61N 1/18 (2006.01)., Method of treatment of patients with gastrointestinal bleeding. U2017 11167, app. 15.11.2017, publ. 25.04.2018 bull №8.
- Chumanevich O.A., Bordakov V.N., Gapanovich V.N. et al. "Endoscopic hemostasis in gastroduodenal bleeding." – Minsk, Medical News Magazine. 2016;8(1).
- Troland D., Stanley A. Endotherapy of Peptic Ulcer Bleeding. Gastrointest Endosc Clin N Am. 2018;28(3):277-289. doi: 10.1016/j.giec.2018.02.002.
- Eun J. G., Ji Y. A., Hwoon-Yong J., Young S. P. et al. Effects of argon plasma coagulation on human stomach tissue: An ex vivo study. J Gastroenterol Hepatol. 2017;32(5):1040-1045. doi: 10.1111/jgh.13636.
- Zippi M., Traversa G., Cocco A. et al. Use of argon plasma coagulation in digestive endoscopy: a concise review. Clin Ter. 2012;163(6):e435-40.
- Gralnek I. M., Dumonceau J-M., Kuipers E. J. et al. Diagnosis and management of nonvariceal upper gastrointestinal hemorrhage: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. Endoscopy. 2015;47(10):a1-46. doi: 10.1055/s-0034-1393172.

ORCID and contributionship:

Yaroslav P. Feleshtynskyi: 0000-0003-4376-4265 ^{A,B,C}
 Sergiy O. Oparin: 0000-0003-1504-9678 ^{A,B,C,F}
 Bogdan V. Sorokin: 0000-0003-0511-7550 ^{D,E}
 Margaryta G. Boiarska: 0000-0002-6853-5533 ^{C, D,E}
 Dmytro V. Lutsenko: 0000-0002-0453-7441 ^{D,E}

Conflict of interest:

The Authors declare no conflict of interest.

CORRESPONDING AUTHOR

Oparin O. Sergiy

Shupyk National Medical Academy of Postgraduate Education of the Ministry of Health of Ukraine
 9 Dorohozhytska St., 04112 Kyiv, Ukraine
 tel: +38(050) 952 46 51
 e-mail: soparin7@gmail.com

Received: 22.04.2021

Accepted: 20.08.2021

A – Work concept and design, B – Data collection and analysis, C – Responsibility for statistical analysis, D – Writing the article, E – Critical review, F – Final approval of the article