

BACTERIAL INFECTION CAUSES OF PREGNANCY LOSS AND PREMATURE BIRTH IN THE WOMEN IN UKRAINE

DOI: 10.36740/WLek202106113

Aidyn G. Salmanov¹, Oleg M. Ishchak^{1,2}, Yuliia M. Shostak¹, Viktoriia V. Kozachenko^{1,3}, Victor O. Rud⁴, Oleg V. Golyanovskiy¹, Volodymyr O. Shkorbotun¹

¹SHUPYK NATIONAL HEALTHCARE UNIVERSITY OF UKRAINE, KYIV, UKRAINE

²TERNOPIL PERINATAL CENTER, TERNOPIL, UKRAINE

³ODESA MATERNITY HOSPITAL, ODESA, UKRAINE

⁴NATIONAL PIROGOV MEMORIAL MEDICAL UNIVERSITY, VINNYTSIA, UKRAINE

ABSTRACT

The aim: To find out whether or not the presence of pregnancy loss and preterm birth was associated with the history of healthcare-associated infection (HAI) of women reproductive tract.

Materials and methods: We performed a retrospective cohort study based on surveillance data. This study included 1,027 fertile women aged 21–50 years admitted to 8 hospitals from 2017–2019 with complaints to pregnancy loss or preterm birth. Definitions HAI of women reproductive tract were used from the CDC/ NHSN.

Results: Of 1,027 fertile women with complaints to pregnancy loss or preterm birth, 702 (68.4%) HAIs of reproductive tract were detected. The predominant HAIs were: postpartum endometritis (19.1%), bacterial vaginitis (15.8%), chorioamnionitis (11.3%), vaginal cuff infection (9.1%), and parametritis (5.6%). According to the statistical data, a significant association between HAI of women reproductive tract and the history of pregnancy loss ($p < 0.05$) was determined. The presence of HAI had no effect on the first trimester miscarriage ($p > 0.05$), but HAI women reproductive tract had strongly affected the second trimester pregnancy losses ($p < 0.05$). Main causative agents of HAI were *Escherichia coli* (31.5%), *Enterobacter* spp. (18.4%), *Klebsiella pneumoniae* (12.5%), *Enterococcus faecalis* (11.6%), *Staphylococcus aureus* (10.2%), *Pseudomonas aeruginosa* (8.4%).

Conclusions: This study identified a significant association between healthcare-associated infection of women reproductive tract and a history of miscarriage. However, there was no association between HAI and a history of preterm birth.

KEY WORDS: pregnancy loss, miscarriage, preterm birth, healthcare-associated infection, reproductive tract infection, bacterial pathogen, antimicrobial resistance

Wiad Lek. 2021;74(6):1355–1359

INTRODUCTION

Pregnancy loss and preterm birth in women is an important public health care concern worldwide. The term reproductive loss is often associated with miscarriage, which manifests itself as spontaneous abortion during the first trimester of pregnancy before 12 weeks (early miscarriage) or 12 to 24 weeks of gestation (late miscarriage) in women [1]. Miscarriage can be also classified as embryonic loss (or early miscarriage) when it occurs before 10 gestational weeks and fetal loss (or fetal miscarriage) when it occurs after 10 gestational weeks [2, 3].

According to literature, research suggests that between 10% and 20% of women with a medically confirmed pregnancy will end in miscarriage [4]. In an Australian prospective cohort women the rate of miscarriage varied from 11.3 to 86.5 per 100 live births; overall, miscarriage occurred in 25% of the women in the study [5]. Early pregnancy loss in women is a relatively common event, occurring in 15%–25% of pregnancies, and increasing in prevalence with maternal age [6]. The risk is between 9% and 12% in women aged ≤ 35 years, but increases to 50% in women aged > 40 . According

to the American College of Obstetricians and Gynecologists, risk of miscarriage in women aged 45 years but increases to 80% [7]. Recurrent pregnancy loss is an important reproductive health issue, affecting 2%–5% of couples [6, 8]. However, almost half of the cases remain unexplained and are empirically treated [8].

Miscarriage is one of the most common reproductive losses, however, the causes and adverse outcomes of pregnancy remain poorly understood. According to literature, common established causes include uterine anomalies [9], age of both parents [5, 6], antiphospholipid syndrome [10], hormonal and metabolic disorders [11], and cytogenetic abnormalities [12]. Other etiologies have been proposed but are still considered controversial, such as chronic endometritis [13, 14] and inherited thrombophilias [15]. In addition, there is evidence in the literature that obesity [6, 16], smoking [17], excessive caffeine consumption [17], and cocaine [18] is associated with an increased risk of miscarriage.

Can be difficult to pinpoint exactly what causes pregnancy loss, there are some infections that are associated with an increased risk of miscarriage. According to literature,

bacterial vaginosis, brucellosis, syphilis, cytomegalovirus, dengue fever, human immunodeficiency virus, rubella, and malaria are more frequently found in women with miscarriage [1]. However, no trials have been published to date, remain concerning the impact of healthcare-associated reproductive tract infection on reproductive outcome.

THE AIM

The aim of this study was to find out whether or not the presence of pregnancy loss and preterm birth was associated with the history of healthcare-associated infection of women reproductive tract.

MATERIALS AND METHODS

SETTING AND PATIENTS

Over a 36-month period (January 2017 to December 2019), this a retrospective cohort study was performed in 8 Ukrainian women's hospitals, which are similar in terms of medical equipment, personnel, laboratory facilities, and number of beds. All participating hospitals were required to have a clinical microbiology laboratory with the capacity to process cultures. This study included fertile women aged 21-50 years admitted to hospitals with complaints to pregnancy loss or preterm birth. All women were local residents. Patients who were referred from other hospitals were excluded. Exclusion criteria were patients with current pregnancy, positive serological test for syphilis or other sexually transmitted infections.

DEFINITIONS

For the purposes of this study miscarriage is defined as the spontaneous loss of a pregnancy during the first 24 weeks of gestation. Diagnosis of healthcare-associated infection (HAI) women reproductive tract was based on criteria from the CDC/NHSN Surveillance Definitions for Specific Types of Infections [19].

DATA COLLECTION

All data on outcome of pregnancy (pregnancy loss and preterm birth) and HAIs of women reproductive tract, and their causative pathogens, were collected retrospectively on a form specifically designed by the investigators using medical records comprising charts, daily flow sheets, and laboratory (microbiology) results.

ETHICS

The Shupyk National Healthcare University of Ukraine Ethics Committee approved the waiver of informed consent to participate in this study due to its retrospective design. All women's data were anonymised prior to the analysis.

MICROBIOLOGICAL METHODS

All samples were obtained from women with clinical symptoms of HAIs. Microbial isolates were identified using

standard microbiological techniques, including automated microbiology testing (Vitek-2; bioMérieux, Marcy l'Etoile, France).

STATISTICAL ANALYSIS

The aim of this study was to find out whether or not the presence of early pregnancy loss or preterm birth was associated with the history of HAI of women reproductive tract. For these comparisons, χ^2 or Fischer's exact test was used, and P values less than 0.05 were considered as statistically significant.

RESULTS

During the study period, 68.4% [95% confidence interval (CI) 66.9 -69.8] (702 of 1,027) fertile women with complaints to pregnancy loss and preterm birth were found to have healthcare-associated reproductive tract infections. Of these cases, 1.7% (17/1027) pelvic abscess or cellulitis, 1.8% (18/1027) infection of the uterine appendages, 2.5% (26/1027) infection of the Fallopian tubes, 1.7% (17/1027) infection of the ovaries, 5.6% (57/1027) infection of the supporting ligaments, 11.3% (116/1027) chorioamnionitis, 15.8% (162/1027) infection of the vagina, 19.1% (196/1027) postpartum endometritis, and 9.1% (93/1027) vaginal cuff infection were identified (Table I).

The women in the study population were also evaluated in view of the miscarriages whether it has happened in the first or second trimester of gestation. 518 (73.8%) of the 702 women had the first trimester miscarriage (≤ 12 weeks), and the remaining 184 (26.2%) of 702 women had the second trimester miscarriage (> 12 weeks).

When we examined the study population with of healthcare-associated infection (HAI), 511 (72.8%) of the 702 women were detected as having the history of a early pregnancy loss in the last 6 months, 191 (27.2%) of 702 women had recurrent pregnancy losses.

In the study population, 511 of 702 (72.8%) women with HAI had a history of early pregnancy loss in the last 6 months and only 191 of 702 (27.2%) women with HAI had recurrent pregnancy losses (Table II). According to the statistical data, a significant association between HAI and the history of pregnancy loss ($P < 0.05$) was determined. However, there was no association between HAI and the history of preterm birth ($P > 0.05$).

In additional, the effect of HAI women reproductive tract on the gestation periods was also examined. The presence of HAI had no effect on the first trimester miscarriage ($P > 0.05$), but HAI women reproductive tract had strongly affected the second trimester pregnancy losses ($P < 0.05$).

In our study, a total of 987 pathogens isolated from 702 cases HAIs of women reproductive tract. Main causative agents of HAI were *Escherichia coli* (31.5%), *Enterobacter* spp. (18.4%), *Klebsiella pneumoniae* (12.5%), *Enterococcus faecalis* (11.6%), *Staphylococcus aureus* (10.2%), *Pseudomonas aeruginosa* (8.4%). The distribution of microorganisms causing HAIs of women reproductive tract is shown in Table III.

Table I. Frequency of pregnancy outcomes in fertile women in Ukraine between 2017 and 2019 by healthcare-associated reproductive tract infection (n = 1027)

HAI ^a type	Number of miscarriages	
	n	%
A pelvic abscess or cellulitis	17	1.7
Infection of the uterine appendages (<i>Adnexa utery</i>)	18	1.8
Infection of the Fallopian tubes (<i>Salpingitis</i>)	26	2.5
Infection of the ovaries (<i>Oophoritis</i>)	17	1.7
Infection of the supporting ligaments (<i>Parametritis</i>)	57	5.6
Chorioamnionitis	116	11.3
Infection of the Vagina (<i>Bacterial Vaginitis</i>)	162	15.8
Postpartum endometritis	196	19.1
Vaginal cuff infection	93	9.1
Total	702	68.4

Note: ^aHAI, healthcare-associated infection

Table II. Relationship among the presence of early pregnancy loss or preterm birth, history of healthcare-associated reproductive tract infection (n=1,027).

Pregnancy outcomes	HAI (+) (n=702, %)	HAI (-) (n=325, %)	P value
Pregnancy loss	511 (72.8)	137 (42.2)	P < 0.05
Preterm birth	191 (27.2)	188 (57.8)	P < 0.05

Table III. The distribution of microorganisms (n = 987) causing HAIs of women reproductive tract

Microorganism	All isolates	
	n	%
<i>Gram-positive cocci</i>	246	24.9
<i>Enterococcus faecalis</i>	114	11.6
<i>Staphylococcus aureus</i>	101	10.2
<i>Streptococcus spp.</i>	14	1.4
<i>Staphylococcus epidermidis</i>	12	1.2
<i>Enterococcus faecium</i>	3	0.3
<i>Other Gram-positive cocci</i>	2	0.2
<i>Gram-negative bacilli</i>	741	75.1
<i>Escherichia coli</i>	313	31.7
<i>Enterobacter spp.</i>	182	18.4
<i>Klebsiella pneumoniae</i>	123	12.5
<i>Pseudomonas aeruginosa</i>	83	8.4
<i>Proteus mirabilis</i>	13	1.3
<i>Serratia spp.</i>	9	0.9
<i>Citrobacter spp.</i>	8	0.8
<i>Klebsiella oxytoca</i>	7	0.7
<i>Other Gram-negative bacilli</i>	3	0.3

DISCUSSION

Although miscarriage is considered the most common adverse pregnancy outcome worldwide, figures for Ukraine are not available. The aim of this study was to find out whether or not the presence of early pregnancy loss and preterm birth was associated with the history of healthcare-associated infection of women reproductive tract. In

our study, HAI women reproductive tract was detected in 68.4% fertile women by criteria from the CDC/NHSN Surveillance Definitions for Specific Types of Infections [19]. Of these cases, 1.7% pelvic abscess or cellulitis, 1.8% infection of the uterine appendages, 2.5% infection of the Fallopian tubes, 1.7% infection of the ovaries, 5.6% infection of the supporting ligaments, 11.3% chorioamnionitis,

15.8% infection of the vagina, 19.1% postpartum endometritis, and 9.1% vaginal cuff infection were identified.

In our study 73.8% women had the first trimester miscarriage, and the remaining 26.2% women had the second trimester miscarriage. When we examined the study population with of HAI, 72.8% women were detected as having the history of a early pregnancy loss in the last 6 months, 27.2% women had recurrent pregnancy losses. In the study population, 72.8% women with HAI had a history of early pregnancy loss in the last 6 months and only 27.2% women with HAI had recurrent pregnancy losses. According to the statistical data, a significant association between HAI and the history of pregnancy loss was determined. However, there was no association between HAI and the history of preterm birth. In additional, the effect of HAI women reproductive tract on the gestation periods was also examined. The presence of HAI had no effect on the first trimester miscarriage, but HAI women reproductive tract had strongly affected the second trimester pregnancy losses.

In our study many women who miscarry do not have any risk factors prior to their miscarriage. In contrast, some women have many risk factors for miscarriage, but end up carrying a normal pregnancy to term.

Reproductive loss is an inclusive term for women, related to reproductive health, fertility, the outcomes of a pregnancy, and includes experiences of miscarriage, stillbirth, infertility, abortion, pregnancy loss, and loss of a child.

According to literature, the definition of recurrent pregnancy loss differs among international societies. For the European Society for Human Reproduction and Embryology [2, 20], and the Royal College of Obstetricians and Gynaecologists [21]. recurrent pregnancy loss refers to three consecutive pregnancy losses, including nonvisualized ones. However, according to the American Society for Reproductive Medicine [6], it is defined as two or more clinical pregnancy losses, but not necessarily consecutive. Early pregnancy loss, also referred to as miscarriage or spontaneous abortion, is defined as the loss of a clinical pregnancy before 20 completed weeks of gestational age (18 weeks after fertilization) or, if gestational age is unknown, the loss of an embryo/fetus of <400 g. [22]. Ectopic, molar, and biochemical pregnancies are thus not included.[6].

We analyzed a large number of bacterial infectious agents to determine if they are associated with an increased risk of miscarriage. Further research is required to clarify whether certain infections do increase miscarriage risk. This issue is of particular importance for public health practitioners as it could alter current policies of prevention of infection, diagnosis and treatment in pregnant women.

CONCLUSIONS

This study identified a significant association between HAI of women reproductive tract and a history of miscarriage. However, there was no association between HAI and a history of preterm birth. For most of the bacterial pathogens the mechanism that leads from infection to miscarriage is unknown. Further research is required to

clarify whether certain reproductive tract infections do increase miscarriage risk.

REFERENCES

1. Giakoumelou S., Wheelhouse N., Cuschieri K. et al. The role of infection in miscarriage. *Hum Reprod Update*. 2016;22(1):116-133. doi:10.1093/humupd/dmv041.
2. Kolte A.M., Bernardi L.A., Christiansen O.B. et al. ESHRE Special Interest Group, Early Pregnancy Terminology for pregnancy loss prior to viability: a consensus statement from the ESHRE early pregnancy special interest group. *Hum Reprod*. 2015;30(3):495-8. doi: 10.1093/humrep/deu299.
3. Silver R.M., Branch D.W., Goldenberg R. et al. Nomenclature for pregnancy outcomes: time for a change. *Obstet Gynecol*. 2011;118(6):1402-8. doi: 10.1097/AOG.0b013e3182392977.
4. Cohain J.S., Buxbaum R.E., Mankuta D. Spontaneous first trimester miscarriage rates per woman among parous women with 1 or more pregnancies of 24 weeks or more. *BMC Pregnancy Childbirth*. 2017;17(1):437. doi:10.1186/s12884-017-1620-1.
5. Hure A.J., Powers J.R., Mishra G.D. et al. Miscarriage, preterm delivery, and stillbirth: large variations in rates within a cohort of Australian women. *PLoS One*. 2012;7(5):e37109. doi: 10.1371/journal.pone.0037109.
6. Practice Committee of the American Society for Reproductive Medicine Evaluation and treatment of recurrent pregnancy loss: a committee opinion. *Fertil Steril*. 2012;98(5):1103-11. doi: 10.1016/j.fertnstert.2012.06.048.
7. American College of Obstetricians and Gynecologists. ACOG practice bulletin No. 200 summary: Early pregnancy loss. *Obstet Gynecol*. 2018;132(5):1311-1313. doi:10.1097/AOG.0000000000002900.
8. El Hachem H., Crepau V., May-Panloup P. et al. Recurrent pregnancy loss: current perspectives. *Int J Womens Health*. 2017;9:331-345. doi:10.2147/IJWH.S100817.
9. Jaslow C.R., Carney J.L., Kutteh W.H. Diagnostic factors identified in 1020 women with two versus three or more recurrent pregnancy losses. *Fertil Steril*. 2010;93(4):1234-43. doi: 10.1016/j.fertnstert.2009.01.166.
10. Andreoli L., Chighizola C.B., Banzato A. et al. Estimated frequency of antiphospholipid antibodies in patients with pregnancy morbidity, stroke, myocardial infarction, and deep vein thrombosis: a critical review of the literature. *Arthritis Care Res (Hoboken)*. 2013;65(11):1869-73. doi: 10.1002/acr.22066.
11. Cleary-Goldman J., Malone F.D., Lambert-Messerlian G. et al. Maternal thyroid hypofunction and pregnancy outcome. *Obstet Gynecol*. 2008;112(1):85-92. doi: 10.1097/AOG.0b013e3181788dd7.
12. Sugiura-Ogasawara M., Ozaki Y., Katano K. et al. Abnormal embryonic karyotype is the most frequent cause of recurrent miscarriage. *Hum Reprod*. 2012;27(8):2297-303. doi: 10.1093/humrep/des179.
13. McQueen D.B., Perfetto C.O., Hazard F.K., Lathi R.B. Pregnancy outcomes in women with chronic endometritis and recurrent pregnancy loss. *Fertil Steril*. 2015;104(4):927-931. doi: 10.1016/j.fertnstert.2015.06.044.
14. Bouet P.E., El Hachem H., Monceau E. et al. Chronic endometritis in women with recurrent pregnancy loss and recurrent implantation failure: prevalence and role of office hysteroscopy and immunohistochemistry in diagnosis. *Fertil Steril*. 2016;105(1):106-10. doi: 10.1016/j.fertnstert.2015.09.025.
15. Lissalde-Lavigne G., Fabbro-Peray P., Cochery-Nouvellon E. et al. Factor V Leiden and prothrombin G20210A polymorphisms as risk factors for miscarriage during a first intended pregnancy: the matched case-control 'NOHA first' study. *J Thromb Haemost*. 2005;3(10):2178-84. doi: 10.1111/j.1538-7836.2005.01581.x.

16. Moussa H.N., Alrais M.A., Leon M.G. et al. Obesity epidemic: impact from preconception to postpartum. *Future Sci OA*. 2016;2(3):FS0137. doi:10.4155/fsoa-2016-0035.
17. Lassi Z.S., Imam A.M., Dean S.V. et al. Preconception care: caffeine, smoking, alcohol, drugs and other environmental chemical/radiation exposure. *Reprod Health*. 2014;11(3):S6. doi: 10.1186/1742-4755-11-S3-S6.
18. Ness R.B., Grisso J.A., Hirshinger N. et al. Cocaine and tobacco use and the risk of spontaneous abortion. *N Engl J Med*. 1999;340(5):333–9. doi: 10.1056/NEJM199902043400501.
19. Horan T.C., Andrus M., Dudeck M.A. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*. 2008;36(5):309–32. doi: 10.1016/j.ajic.2008.03.002.
20. Jauniaux E., Farquharson R.G., Christiansen O.B. et al. Evidence-based guidelines for the investigation and medical treatment of recurrent miscarriage. *Hum Reprod*. 2006;21(9):2216–22. doi: 10.1093/humrep/del150.
21. Royal College of Obstetricians and Gynaecologists, Scientific Advisory Committee, Guideline No. 17. The Investigation and treatment of couples with recurrent miscarriage. 2011. <http://www.rcog.org.uk/womens-health/clinical-guidance/investigation-and-treatment-couples-recurrent-miscarriage-green-top->.
22. Zegers-Hochschild F., Adamson G.D., de Mouzon J. et al. International Committee for Monitoring Assisted Reproductive Technology; World Health Organization. International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary of ART terminology, 2009. *Fertil Steril*. 2009;92(5):1520–4. doi: 10.1016/j.fertnstert.2009.09.009.

We would like to thank all the nurses and physicians who contributed to the prevalence surveys.

This work is a fragment of a research study of the Scientific Research Laboratory of Shupyk National Healthcare University of Ukraine (Kyiv, Ukraine). Title: The scientific justification for measures to combat the resistance of microorganisms to antimicrobial drugs in Ukraine on the One Health approaches. State Registration Number: 0120U101440. Study period: 2020-2022. This work was funded by the Ministry of Health of Ukraine according to the plan of that research study. The authors did not receive any financial support from the manufacturers of medical instruments and drugs.

ORCID and contributionship:

Aidyn G. Salmanov: 0000-0002-4673-1154 ^{A, C, D, E, F}

Oleg M. Ishchak: 0000 0002 9634 3635 ^{B, C, D, F}

Yuliia M. Shostak: 0000-0003-2910-5028, ^{B, C, D, F}

Viktoriiia V. Kozachenko: 0000-0002-4140-0663 ^{B, C, D, F}

Victor O. Rud: 0000-0002-0768-6477 ^{B, C, D, F}

Oleg V. Golyanovskiy: 0000-0002-5524-4411 ^{C, D, F}

Volodymyr O. Shkorbotun: 0000-0003-0769-6242 ^{B, C, F}

Conflict of interest:

The Authors declare no conflict of interest

CORRESPONDING AUTHOR

Aidyn G. Salmanov

Shupyk National Healthcare University of Ukraine

9 Dorohozhytska St., 04112 Kyiv, Ukraine

tel: +380667997631

e-mail: mozsago@gmail.com

Received: 05.12.2020

Accepted: 27.04.2021

A - Work concept and design, **B** – Data collection and analysis, **C** – Responsibility for statistical analysis, **D** – Writing the article, **E** – Critical review, **F** – Final approval of the article